

Medical Drug Clinical Criteria

Subject:	Halaven (eribulin)		
Document #:	CC-0108	Publish Date:	03/27/2023
Status:	Reviewed	Last Review Date:	02/24/2023

Table of Contents

Overview	Coding	References
Clinical criteria	Document history	

Overview

This document addresses the use of Halaven (eribulin). Halaven is a non-taxane microtubule dynamics inhibitor that is a synthetic analogue of halichondrin B, a product isolated from a marine sponge. Although the exact mechanism is unknown, it is believed to work through inhibition of the growth phase of microtubule dynamics, without affecting the shortening phase, sequestering tubulin into nonproductive aggregates.

The FDA approved indications for Halaven include metastatic breast cancer or unresectable or metastatic liposarcoma. The National Comprehensive Cancer Network (NCCN) provides additional recommendations with a category 1 and 2A level of evidence for the uses in invasive breast cancer and soft tissue sarcoma.

Definitions and Measures

Adjuvant therapy: Treatment given after the primary treatment to increase the chances of a cure; may include chemotherapy, radiation, hormone or biological therapy.

Anthracycline: A type of antibiotic that comes from certain types of Streptomyces bacteria and are used to treat many types of cancer. Anthracyclines damage the DNA in cancer cells, causing the cells to die.

Gleason Grading system: A prostate cancer grading system. A primary and secondary pattern, the number range of each is from 1 to 5, are assigned and then summed to yield a total score.

Human epidermal growth factor 2 (ERBB2) status: A laboratory finding related to the presence or absence of cellular receptors for HER2/neu; also known as ErbB-2 protein family.

Line of Therapy:

- **First-line therapy:** The first or primary treatment for the diagnosis, which may include surgery, chemotherapy, radiation therapy or a combination of these therapies.
- **Second-line therapy:** Treatment given when initial treatment (first-line therapy) is not effective or there is disease progression.
- **Third-line therapy:** Treatment given when both initial (first-line therapy) and subsequent treatment (second-line therapy) are not effective or there is disease progression.

Metastasis: The spread of cancer from one part of the body to another; a metastatic tumor contains cells that are like those in the original (primary) tumor and have spread.

Microtubule inhibitors (MTI): A class of drugs including taxanes, vinca alkaloids, and epothilones that stabilize or destabilize microtubules, thereby suppressing microtubule dynamics required for proper mitotic function, effectively blocking cell cycle progression and resulting in cell death.

Relapse or recurrence: After a period of improvement, during which time a disease (for example, cancer) could not be detected, the return of signs and symptoms of illness or disease. For cancer, it may come back to the same place as the original (primary) tumor or to another place in the body.

One line of therapy: Single line of therapy.

Taxane: A type of mitotic inhibitor and antimicrotubule drug used to treat cancer that blocks cell growth by stopping mitosis (cell division).

Clinical Criteria

When a drug is being reviewed for coverage under a member's medical benefit plan or is otherwise subject to clinical review (including prior authorization), the following criteria will be used to determine whether the drug meets any applicable medical necessity requirements for the intended/prescribed purpose.

Halaven (eribulin mesylate)

Requests for Halaven (eribulin mesylate) may be approved if the following criteria are met:

- I. Individual has a diagnosis of locally recurrent or metastatic breast cancer (Label, NCCN 2A); **AND**
 - II. Individual is using as monotherapy; **AND**
 - III. Individual is using as a single line of therapy; **AND**
 - IV. Individual has previously received at least two chemotherapeutic regimens for locally recurrent or metastatic disease;
- OR**
- V. Individual has a diagnosis of locally recurrent or metastatic HER2 positive breast cancer (NCCN 2A); **AND**
 - VI. In one of the following ways:
 - A. Individual is using in combination with trastuzumab (or trastuzumab biosimilars); **OR**
 - B. Individual is using in combination with Margenza (margetuximab-cmkb) as third line therapy;
- AND**
- VII. Individual has symptomatic visceral disease; **OR**
 - VIII. Individual has either hormone receptor-negative disease or hormone-receptor positive and endocrine refractory disease;
- OR**
- IX. Individual has a diagnosis of locally recurrent or metastatic soft tissue sarcoma (Label, NCCN 1, 2A); **AND**
 - X. Individual is using as a monotherapy; **AND**
 - XI. Individual is using as a single line of therapy; **AND**
 - XII. Individual has previously received at least two chemotherapeutic regimens for locally recurrent or metastatic disease.

Requests for Halaven may not be approved for the following criteria:

- I. Individual has a diagnosis of head and neck cancer; **OR**
- II. Individual has a diagnosis of non-small cell lung cancer; **OR**
- III. When the above criteria are not met and for all other indications.

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

HCPCS

J9179 Injection, eribulin mesylate, 0.1 mg

ICD-10 Diagnosis

C48.0-C48.8 Malignant neoplasm of retroperitoneum and peritoneum
C49.0-C49.9 Malignant neoplasm of other connective and soft tissue
C50.011-C50.929 Malignant neoplasm of breast
C79.81 Secondary malignant neoplasm of breast
Z17.0 Estrogen receptor negative status [ER-]
Z17.1 Estrogen receptor positive status [ER+]

Document History

Reviewed: 02/24/2023

Document History:

- 02/24/2023 – Annual Review: No Changes. Coding Reviewed: No changes.

- 02/25/2022 – Annual Review: Update breast cancer criteria to include NCCN 2A recommendation use with Margenza in invasive breast cancer. Coding Reviewed: No changes.
- 02/19/2021 – Annual Review: Clarify criteria reference with NCCN 2A notation. Coding Reviewed: No changes.
- 02/21/2020 – Annual Review: Clarify clinical criteria for trastuzumab or trastuzumab biosimilars. Add may not be approved criteria. Coding review: No changes
- 11/15/2019 – Annual Review: No changes. Coding reviewed: No changes
- 05/17/2019 – Annual Review: Initial review of Halaven (eribulin mesylate). Minor wording and formatting updates. Coding Reviewed: Added Z17.0-Z17.1 Hormone receptor status.

References

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2023. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. Cortes J, O'Shaughnessy J, Loesch D, et al. Eribulin monotherapy versus treatment of physician's choice in patients with metastatic breast cancer (EMBRACE): a phase 3 open-label randomised study. *Lancet*. 2011;377(9769):914-923. doi:10.1016/S0140-6736(11)60070-6 Available at: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(11\)60070-6/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(11)60070-6/fulltext). Accessed January 17, 2023
3. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Updated periodically.
4. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
5. Kaufman PA, Awada A, Twelves C, et al. Phase III open-label randomized study of eribulin mesylate versus capecitabine in patients with locally advanced or metastatic breast cancer previously treated with an anthracycline and a taxane. *J Clin Oncol*. 2015;33(6):594-601. doi:10.1200/JCO.2013.52.4892. Available at: <https://ascopubs.org/doi/full/10.1200/JCO.2013.52.4892>. Accessed January 17, 2023
6. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2023; Updated periodically.
7. NCCN Clinical Practice Guidelines in Oncology™. © 2023 National Comprehensive Cancer Network, Inc. For additional information visit the NCCN website: <http://www.nccn.org/index.asp>. Accessed on January 17, 2023.
 - a. Breast Cancer. V4.2022. Revised June 21, 2022.
 - b. Soft Tissue Sarcoma. V2.2022. Revised May 17, 2022.

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

© CPT Only – American Medical Association